

The Contragredient

Jeffrey Adams and David A. Vogan Jr.

January 2, 2012

1 Introduction

Perhaps it is surprising that the following question has not been addressed in the literature: what is the contragredient in terms of Langlands parameters?

Thus G is a connected, reductive algebraic group defined over a local field F , and $G(F)$ is its F -points. According to the local Langlands conjecture, associated to a homomorphism ϕ from the Weil-Deligne group of F into the L -group of $G(F)$ is an L -packet $\Pi(\phi)$, a finite set of irreducible admissible representations of $G(F)$. Conjecturally these sets partition the admissible dual.

So suppose π is an irreducible admissible representation, and $\pi \in \Pi(\phi)$. Let π^* be the contragredient of π . The question is: what is the homomorphism ϕ^* such that $\pi^* \in \Pi(\phi^*)$? We also consider the related question of describing the *Hermitian dual* in terms of Langlands parameters.

Let ${}^\vee G$ be the complex dual group of G . The Chevalley involution C of ${}^\vee G$ satisfies $C(h) = h^{-1}$, for all h in some Cartan subgroup of ${}^\vee G$. The L -group ${}^L G$ of $G(F)$ is a certain semidirect product ${}^\vee G \rtimes \Gamma$ where Γ is the absolute Galois group of F (or other related groups). We can choose C so that it extends to an involution of ${}^L G$, acting trivially on Γ . We refer to this as the Chevalley involution of ${}^L G$. See Section 2.

We believe the contragredient should correspond to composition with the Chevalley involution of ${}^L G$. To avoid two levels of conjecture, we formulate this as follows.

Conjecture 1.1 *Assume the local Langlands conjecture is known for both π and π^* . Let C be the Chevalley involution of ${}^L G$. Then*

$$\pi \in \Pi(\phi) \Leftrightarrow \pi^* \in \Pi(C \circ \phi).$$

Even the following weaker result is not known:

Conjecture 1.2 *If Π is an L-packet, then so is $\{\pi^* \mid \pi \in \Pi\}$.*

The local Langlands conjecture is only known, for fixed $G(F)$ and all π , in a limited number of cases, notably $GL(n, F)$ over any local field, and for any G if $F = \mathbb{R}$ or \mathbb{C} . See Langlands's original paper [10], which is summarized in Borel's article [6]. On the other hand it is known for a restricted class of representations for more groups, for example unramified principal series representations of a split p-adic group [6, 10.4].

It would be reasonable to impose Conjecture 1.1 as a condition on the local Langlands correspondence in cases where it is not known.

We concentrate on the Archimedean case. Let $W_{\mathbb{R}}$ be the Weil group of \mathbb{R} . The contragredient in this case can be realized either via the Chevalley automorphism of ${}^L G$, or via a similar automorphism of $W_{\mathbb{R}}$. By analogy with C , there is a unique \mathbb{C}^* -conjugacy class of automorphisms τ of $W_{\mathbb{R}}$ satisfying $\tau(z) = z^{-1}$ for all $z \in \mathbb{C}^*$. See Section 2.

Theorem 1.3 *Let $G(\mathbb{R})$ be the real points of a connected reductive algebraic group defined over \mathbb{R} , with L-group ${}^L G$. Suppose $\phi : W_{\mathbb{R}} \rightarrow {}^L G$ is an admissible homomorphism, with associated L-packet $\Pi(\phi)$. Let $\Pi(\phi)^* = \{\pi^* \mid \pi \in \Pi(\phi)\}$. Then*

$$\Pi(\phi)^* = \Pi(C \circ \phi) = \Pi(\phi \circ \tau).$$

In particular $\Pi(\phi)^$ is an L-packet.*

Here is a sketch of the proof.

It is easy to prove in the case of tori. See Section 3.

It is well known that an L-packet Π of (relative) discrete series representations is determined by an infinitesimal and a central character. In fact something stronger is true. Let G_{rad} be the radical of G , i.e. the maximal central torus. Then Π is determined by an infinitesimal character and a character of $G_{\text{rad}}(\mathbb{R})$, which we refer to as a radical character.

In general it is easy to read off the infinitesimal and radical characters of $\Pi(\phi)$ [6]. In particular for a relative discrete series parameter the Theorem reduces to a claim about how C affects the infinitesimal and radical characters. For the radical character this reduces to the case of tori, and Theorem 1.3 follows in this case.

This is the heart of the matter, and the general case follows easily by parabolic induction. In other words, the proof relies on the fact that the parameterization is uniquely characterized by:

1. Infinitesimal character,
2. Radical character,
3. Compatibility with parabolic induction.

For the convenience of the reader we have included a self-contained description of the local Langlands classification in these terms. We make extensive use of the Tits group (see Section 5), which simplifies some technical arguments.

Now consider $GL(n, F)$ for F a local field of characteristic 0. Since $GL(n, F)$ is split we may take ${}^L G = GL(n, \mathbb{C})$ (i.e. $\Gamma = 1$), and an admissible homomorphism ϕ is an n -dimensional representation of the Weil-Deligne group W'_F . In this case L-packets are singletons, so write $\pi(\phi)$ for the representation attached to ϕ .

For the Chevalley involution take $C(g) = {}^t g^{-1}$. Then $C \circ \phi \simeq \phi^*$, the contragredient of ϕ . Over \mathbb{R} Theorem 1.3 says the Langlands correspondence commutes with the contragredient:

$$(1.4) \quad \pi(\phi^*) \simeq \pi(\phi)^*.$$

This is also true over a p-adic field [7], [8], in which case it is closely related to the functional equations for L and ε factors.

We now consider a variant of (1.4) in the real case. Suppose π is an irreducible representation of $GL(n, \mathbb{R})$. Its Hermitian dual π^h is a certain irreducible representation, such that $\pi \simeq \pi^h$ if and only if the space of π supports an invariant Hermitian form. We say π is Hermitian if $\pi \simeq \pi^h$. See Section 8.

The Hermitian dual arises naturally in the study of unitary representations: the unitary dual is the subset of the fixed points of this involution,

consisting of those π for which the invariant form is definite. So it is natural to ask what the Hermitian dual is on the level of Langlands parameters.

There is a natural notion of Hermitian dual of a finite dimensional representation ϕ of $W_{\mathbb{R}}$: $\phi^h = {}^t\bar{\phi}^{-1}$, and ϕ preserves a nondegenerate Hermitian form if and only if $\phi \simeq \phi^h$.

The local Langlands correspondence for $GL(n, \mathbb{R})$ commutes with the Hermitian dual operation:

Theorem 1.5 *Suppose ϕ is an n -dimensional semisimple representation of $W_{\mathbb{R}}$. Let $\phi^h = {}^t\bar{\phi}^{-1}$ be its Hermitian dual. Then:*

1. $\pi(\phi^h) = \pi(\phi)^h$,
2. ϕ is Hermitian if and only if $\pi(\phi)$ is Hermitian,
3. ϕ is unitary if and only if $\pi(\phi)$ is tempered.

See Section 8.

Return now to the setting of general real groups. The space \mathcal{X}_0 of conjugacy classes of L-homomorphisms parametrizes L-packets of representations. By introducing some extra data we obtain a space \mathcal{X} which parametrizes irreducible representations [3]. Roughly speaking \mathcal{X} is the set of conjugacy classes of pairs (ϕ, χ) where $\phi \in \mathcal{X}_0$ and χ is a character of the component group of $\text{Cent}_{\vee G}(\phi(W_{\mathbb{R}}))$. It is natural to ask for the involution of \mathcal{X} induced by the contragredient.

On the other hand, it is possible to formulate and prove an analogue of Theorem 1.5 for general real groups, in terms of an anti-holomorphic involution of ${}^L G$. Also, the analogue of Theorem 1.5 holds in the p-adic case. All of these topics require more machinery. In an effort to keep the presentation as elementary as possible we defer them to a later paper.

We thank Kevin Buzzard for asking about the contragredient on the level of L-parameters.

2 The Chevalley Involution

We discuss the Chevalley involution. This is well known, but for the convenience of the reader we give complete details. We also discuss a similar involution of $W_{\mathbb{R}}$.

Throughout this paper G is a connected, reductive algebraic group. We identify it with its complex points, and write $G(\mathbb{C})$ on occasion to emphasize this point of view. For $x \in G$ write $\text{int}(x)$ for the inner automorphism $\text{int}(x)(g) = xgx^{-1}$.

Proposition 2.1 *Fix a Cartan subgroup H of G . There is an automorphism C of G satisfying $C(h) = h^{-1}$ for all $h \in H$. For any such automorphism $C^2 = 1$, and, for every semisimple element g , $C(g)$ is conjugate to g^{-1} .*

Suppose C_1, C_2 are two such automorphisms defined with respect to Cartan subgroups H_1 and H_2 . Then C_1 and C_2 are conjugate by an inner automorphism of G .

The proof uses *based root data* and *pinning*s. For background see [12]. Fix a Borel subgroup B of G , and a Cartan subgroup $H \subset B$. Let $X^*(H), X_*(H)$ be the character and co-character lattices of H , respectively. Let $\Pi, \vee\Pi$ be the sets of simple roots, respectively simple co-roots, defined by B .

The based root datum defined by (B, H) is $(X^*(H), \Pi, X_*(H), \vee\Pi)$. There is a natural notion of isomorphism of based root data. A pinning is a set $\mathcal{P} = (B, H, \{X_\alpha | \alpha \in \Pi\})$ where, for each $\alpha \in \Pi$, X_α is contained in the α -root space \mathfrak{g}_α of $\mathfrak{g} = \text{Lie}(G)$. Let $\text{Aut}(\mathcal{P})$ be the subgroup of $\text{Aut}(G)$ preserving \mathcal{P} . We refer to the elements of $\text{Aut}(\mathcal{P})$ as \mathcal{P} -distinguished automorphisms.

Theorem 2.2 ([12]) *Suppose G, G' are connected, reductive complex groups. Fix pinning*s $(B, H, \{X_\alpha\})$ and $\mathcal{P} = (B', H', \{X'_\alpha\})$. Let D_b, D'_b be the based root data defined by (B, H) and (B', H') .

Suppose $\phi : D_b \rightarrow D'_b$ is an isomorphism of based root data. Then there is a unique isomorphism $\psi : G \rightarrow G'$ taking \mathcal{P} to \mathcal{P}' and inducing ϕ on the root data.

The only inner automorphism in $\text{Aut}(\mathcal{P})$ is the identity, and there are isomorphisms

$$(2.3) \quad \text{Out}(G) \simeq \text{Aut}(D_b) \simeq \text{Aut}(\mathcal{P}) \subset \text{Aut}(G).$$

The following consequence of the Theorem is quite useful.

Lemma 2.4 *Suppose $\tau \in \text{Aut}(G)$ restricts trivially to a Cartan subgroup H . Then $\tau = \text{int}(h)$ for some $h \in H$.*

Proof. Fix a pinning $(B, H, \{X_\alpha\})$. Then $d\tau(\mathfrak{g}_\alpha) = \mathfrak{g}_\alpha$ for all α . Therefore we can choose $h \in H$ so that $d\tau(X_\alpha) = \text{Ad}(h)(X_\alpha)$ for all $\alpha \in \Pi$. Then $\tau \circ \text{int}(h^{-1})$ acts trivially on D_b and \mathcal{P} . By the theorem $\tau = \text{int}(h)$. \square

Proof of the Proposition. Choose a Borel subgroup B containing H and let $D_b = (X^*(H), \Pi, X_*(H), \vee\Pi)$ be the based root datum defined by (B, H) . Let B^{op} be the opposite Borel, with corresponding root datum $D_b^{op} = (X^*(H), -\Pi, X_*(H), -\vee\Pi)$.

Choose a pinning $\mathcal{P} = (B, H, \{X_\alpha\})$. Let $\mathcal{P}^{op} = (H, B^{op}, \{X_{-\alpha} | \alpha \in \Pi\})$ where, for $\alpha \in \Pi$, $X_{-\alpha} \in \mathfrak{g}_{-\alpha}$ satisfies $[X_\alpha, X_{-\alpha}] = \vee\alpha$.

Let $\phi : D_b \rightarrow D_b'$ be the isomorphism of based root data given by -1 on $X^*(H)$. By the Theorem there is an automorphism $C_{\mathcal{P}}$ of G taking \mathcal{P} to \mathcal{P}^{op} and inducing ϕ . In particular $C_{\mathcal{P}}(h) = h^{-1}$ for $h \in H$. This implies $C_{\mathcal{P}}(\mathfrak{g}_\alpha) = \mathfrak{g}_{-\alpha}$, and since $C_{\mathcal{P}} : \mathcal{P} \rightarrow \mathcal{P}^{op}$ we have $C_{\mathcal{P}}(X_\alpha) = X_{-\alpha}$. Since $C_{\mathcal{P}}^2$ is an automorphism of G , taking \mathcal{P} to itself, and inducing the trivial automorphism of D_b the Theorem implies $C_{\mathcal{P}}^2 = 1$.

If g is any semisimple element, choose x so that $xgx^{-1} \in H$. Then $C(g) = C(x^{-1}(xgx^{-1})x) = (C(x^{-1})x)g^{-1}(C(x^{-1})x)^{-1}$.

Suppose $C_1(h) = h^{-1}$ for all $h \in H$. Then $C_1 \circ C_{\mathcal{P}}$ acts trivially on H , so by the Lemma $C_1 = \text{int}(h) \circ C_{\mathcal{P}}$, which implies $C_1^2 = 1$.

For the final assertion choose g so that $gH_1g^{-1} = H_2$. Then $\text{int}(g) \circ C_1 \circ \text{int}(g^{-1})$ acts by inversion on H_2 . By the Lemma $\text{int}(g) \circ C_1 \circ \text{int}(g^{-1}) = \text{int}(h_2) \circ C_2$ for some $h_2 \in H_2$. Choose $t \in H_2$ so that $t^2 = h_2$. Then $\text{int}(t^{-1}g) \circ C_1 \circ \text{int}(t^{-1}g)^{-1} = C_2$. \square

An involution satisfying the condition of the Proposition is known as a *Chevalley involution*. For \mathcal{P} a splitting datum we refer to the involution $C_{\mathcal{P}}$ of the proof as the *Chevalley involution defined by \mathcal{P}* . The proof shows that every Chevalley involution is equal to $C_{\mathcal{P}}$ for some \mathcal{P} , and all Chevalley involutions are conjugate. We will abuse notation slightly and refer to *the* Chevalley involution.

Remark 2.5

1. If $G = GL(n)$, $C(g) = {}^t g^{-1}$ is a Chevalley involution. Then $G^C = O(n, \mathbb{C})$, the complexified maximal compact subgroup of $GL(n, \mathbb{R})$. In other words, C is the Cartan involution for $GL(n, \mathbb{R})$. In general the Chevalley involution is the Cartan involution of the split real form of G .
2. Suppose C' is any automorphism such that $C'(g)$ is G -conjugate to g^{-1} for all semisimple g . It is not hard to see, using Lemma 2.4, that

$C' = \text{int}(g) \circ C$ for some $g \in G$ and some Chevalley involution C . In other words these automorphisms determine a distinguished element of $\text{Out}(G)$.

3. The Chevalley involution is inner if and only if G is semisimple and -1 is in the Weyl group, in which case $C = \text{int}(g_0)$ where $g_0 \in \text{Norm}_G(H)$ represents w_0 . The Proposition implies g_0^2 is central, and independent of all choices. See Lemma 5.4.

Lemma 2.6 *Fix a splitting datum \mathcal{P} . Then $C_{\mathcal{P}}$ commutes with every \mathcal{P} -distinguished automorphism.*

This is immediate.

Here is a similar involution of $W_{\mathbb{R}}$. Recall $W_{\mathbb{R}} = \langle \mathbb{C}^*, j \rangle$ with relations $jzj^{-1} = \bar{z}$ and $j^2 = -1$.

Lemma 2.7 *There is an involution τ of $W_{\mathbb{R}}$ such that $\tau(z) = z^{-1}$ for all $z \in \mathbb{C}^*$. Any two such automorphisms are conjugate by $\text{int}(z)$ for some $z \in \mathbb{C}^*$.*

Proof. This is elementary. For $z_0 \in \mathbb{C}^*$ define $\tau_{z_0}(z) = z^{-1}$ ($z \in \mathbb{C}^*$) and $\tau_{z_0}(j) = z_0j$. From the relations this extends to an automorphism of $W_{\mathbb{R}}$ if and only if $z_0\bar{z}_0 = 1$. Thus τ_1 is an automorphism, and $\tau_{z_0} = \text{int}(u) \circ \tau_1 \circ \text{int}(u^{-1})$, provided $(u/\bar{u})^2 = z_0$. \square

We refer to such an involution as the Chevalley involution of $W_{\mathbb{R}}$.

3 Tori

Let H be a complex torus, and fix an element $\gamma \in \frac{1}{2}X^*(H)$. Let

$$(3.1) \quad H_{\gamma} = \{(h, z) \in H \times \mathbb{C}^* \mid 2\gamma(h) = z^2\}.$$

This is a two-fold cover of H via the map $(h, z) \rightarrow h$; write ζ for the nontrivial element in the kernel of this map. We call this the γ -cover of H . Note that $(h, z) \rightarrow z$ is a genuine character of H_{γ} , and is a canonical square root of 2γ , denoted γ .

Now assume H is defined over \mathbb{R} , with Cartan involution θ . The γ cover of $H(\mathbb{R})$ is defined to be the inverse image of $H(\mathbb{R})$ in H_{γ} . A character of $H(\mathbb{R})_{\gamma}$ is said to be genuine if it is nontrivial on ζ .

Lemma 3.2 ([5], Proposition 5.8) *Given $\gamma \in \frac{1}{2}X^*(H)$, the genuine characters of $H(\mathbb{R})_\gamma$ are canonically parametrized by the set of pairs (λ, κ) with $\lambda \in \mathfrak{h}^*$, $\kappa \in \gamma + X^*(H)/(1 - \theta)X^*(H)$, and satisfying $(1 + \theta)\lambda = (1 + \theta)\kappa$.*

Write $\chi(\lambda, \kappa)$ for the character defined by (λ, κ) . This character has differential λ , and its restriction to the maximal compact subgroup is the restriction of the character κ of H_γ .

Let ${}^\vee H$ be the dual torus. This satisfies: $X^*({}^\vee H) = X_*(H)$, $X_*({}^\vee H) = X^*(H)$. If H is defined over \mathbb{R} , with Cartan involution θ , then θ may be viewed as an involution of $X_*(H)$; its adjoint θ^t is an involution of $X^*(H) = X_*({}^\vee H)$. Let ${}^\vee\theta$ be the automorphism of ${}^\vee H$ induced by $-\theta^t$.

The L-group of H is ${}^L H = \langle {}^\vee H, {}^\vee\delta \rangle$ where ${}^\vee\delta^2 = 1$ and ${}^\vee\delta$ acts on ${}^\vee H$ by ${}^\vee\theta$. Part of the data is the distinguished element ${}^\vee\delta$ (more precisely its conjugacy class).

More generally an E-group for H is a group ${}^E H = \langle {}^\vee H, {}^\vee\delta \rangle$, where ${}^\vee\delta$ acts on ${}^\vee H$ by ${}^\vee\theta$, and ${}^\vee\delta^2$ is allowed to be an arbitrary element of ${}^\vee H^{\vee\theta}$. Such a group is determined up to isomorphism by the image of ${}^\vee\delta^2$ in ${}^\vee H^{\vee\theta}/\{h{}^\vee\theta(h) \mid h \in {}^\vee H\}$. Again the data includes the ${}^\vee H$ conjugacy class of ${}^\vee\delta$. See [5, Definition 5.9].

A homomorphism $\phi : W_\mathbb{R} \rightarrow {}^E H$ is said to be admissible if it is continuous and $\phi(j) \in {}^L H \setminus {}^\vee H$. Admissible homomorphisms parametrize representations of the covers $H(\mathbb{R})_\gamma$.

Lemma 3.3 ([5], Theorem 5.11) *In the setting of Lemma 3.2, suppose $(1 - \theta)\gamma \in X^*(H)$. View γ as an element of $\frac{1}{2}X_*({}^\vee H)$. Let ${}^E H = \langle {}^\vee H, {}^\vee\delta \rangle$ where ${}^\vee\delta$ acts on ${}^\vee H$ by ${}^\vee\theta$, and ${}^\vee\delta^2 = \exp(2\pi i\gamma) \in {}^\vee H^{\vee\theta}$.*

There is a canonical bijection between the irreducible genuine characters of $H(\mathbb{R})_\gamma$ and ${}^\vee H$ -conjugacy classes of admissible homomorphisms $\phi : W_\mathbb{R} \rightarrow {}^L H$.

Sketch of proof. If ϕ is an admissible homomorphism it may be written in the form

$$(3.4) \quad \begin{aligned} \phi(z) &= z^\lambda \bar{z}^{\vee\theta(\lambda)} \\ \phi(j) &= \exp(2\pi i\mu)^{\vee\delta} \end{aligned}$$

for some $\lambda, \mu \in \vee\mathfrak{h}$. Then $\phi(j)^2 = \exp(2\pi i(\mu + \vee\theta\mu) + \gamma)$ and $\phi(-1) =$

$\exp(\pi i(\lambda - \vee\theta\lambda))$, so $\phi(j^2) = \phi(j)^2$ if and only if

$$(3.5) \quad \kappa := \frac{1}{2}(1 - \vee\theta)\lambda - (1 + \vee\theta)\mu \in \gamma + X_*(\vee H) = \gamma + X^*(H).$$

Then $(1 + \theta)\lambda = (1 + \theta)\kappa$; take ϕ to $\chi(\lambda, \kappa)$. □

Write $\chi(\phi)$ for the genuine character of $H(\mathbb{R})_\gamma$ associated to ϕ .

The Chevalley involution C of $\vee H$ (i.e. inversion) extends to an involution of ${}^E H = \langle \vee H, \vee\delta \rangle$, fixing $\vee\delta$ (this uses the fact that $\exp(2\pi i(2\gamma)) = 1$).

Here is the main result in the case of (covers of) tori.

Lemma 3.6 *Suppose $\phi : W_{\mathbb{R}} \rightarrow {}^E H$ is an admissible homomorphism, with corresponding genuine character $\chi(\phi)$ of $H(\mathbb{R})_\gamma$. Then*

$$(3.7) \quad \chi(C \circ \phi) = \chi(\phi)^*$$

Proof. Suppose ϕ is given by (6.4), so $\chi(\phi) = \chi(\lambda, \kappa)$ with κ as in (3.5). Then

$$(3.8) \quad \begin{aligned} (C \circ \phi)(z) &= z^{-\lambda} \bar{z}^{-\vee\theta(\lambda)} \\ (C \circ \phi)(j) &= \exp(-2\pi i \mu)^{\vee\delta} \end{aligned}$$

By (3.5) $\chi(C \circ \phi) = \chi(-\lambda, -\kappa) = \chi(\lambda, \kappa)^*$. □

4 L-packets without L-groups

We describe L-packets in terms of data for G itself. For the relation with L-parameters see Section 6.

Suppose G is defined over \mathbb{R} , with real form $G(\mathbb{R})$. Thus $G(\mathbb{R}) = G(\mathbb{C})^\sigma$ where σ is an antiholomorphic involution. Fix a Cartan involution θ of G , and let $K = G^\theta$. By definition, $K \cap G(\mathbb{R})$ is a maximal compact subgroup of $G(\mathbb{R})$, with complexification K .

Suppose H is a θ -stable Cartan subgroup of G . After conjugating by K we may assume it is defined over \mathbb{R} , which we always do without further comment.

The imaginary roots Δ_i , i.e. those fixed by θ , form a root system. Let ρ_i be one-half the sum of a set Δ_i^+ of positive imaginary roots. The two-fold

cover H_{ρ_i} of H is defined as in Section 3. It is convenient to eliminate the dependence on Δ_i^+ : define \widetilde{H} to be the inverse limit of $\{H_{\rho_i}\}$ over all choices of Δ_i^+ . The inverse image of $H(\mathbb{R})$ in H_{ρ_i} is denoted $H(\mathbb{R})_{\rho_i}$, and take the inverse limit to define $\widetilde{H(\mathbb{R})}$.

Definition 4.1 *An L-datum is a pair (H, Λ) where H is a θ -stable Cartan subgroup of G , Λ is a genuine character of $\widetilde{H(\mathbb{R})}$, and $\langle d\Lambda, \vee\alpha \rangle \neq 0$ for all imaginary roots.*

Associated to each L-datum is an L-packet. We start by defining relative discrete series L-packets.

We say $H(\mathbb{R})$ is *relatively compact* if $H(\mathbb{R}) \cap G_d$ is compact, where G_d is the derived group of G . Then $G(\mathbb{R})$ has relative discrete series representations if and only if it has a relatively compact Cartan subgroup.

Suppose $H(\mathbb{R})$ is relatively compact. Choose a set of positive roots Δ^+ and define the Weyl denominator

$$(4.2) \quad D(\Delta^+, h) = \prod_{\alpha \in \Delta^+} (1 - e^{-\alpha(h)}) e^{\rho(h)} \quad (h \in H(\mathbb{R})_{\rho}).$$

This is a genuine function, i.e. satisfies $D(\Delta^+, \zeta h) = -D(\Delta^+, h)$. We may view this as a function on $\widetilde{H(\mathbb{R})}$.

Let $q = \frac{1}{2} \dim(G_d/K \cap G_d)$. Let $W(K, H) = \text{Norm}_K(H)/H \cap K$; this is isomorphic to the real Weyl group $W(G(\mathbb{R}), H(\mathbb{R})) = \text{Norm}_{G(\mathbb{R})}(H(\mathbb{R}))/H(\mathbb{R})$.

Definition 4.3 *Suppose $\gamma = (H, \Lambda)$ is an L-datum with $H(\mathbb{R})$ relatively compact. Let $\pi = \pi(\gamma)$ be the unique, non-zero, relative discrete series representation whose character restricted to the regular elements of $H(\mathbb{R})$ is*

$$(4.4) \quad \Theta_{\pi}(h) = (-1)^q D(\Delta^+, \tilde{h})^{-1} \sum_{w \in W(K, H)} \text{sgn}(w)(w\Lambda)(\tilde{h})$$

where $\tilde{h} \in \widetilde{H(\mathbb{R})}$ is any inverse image of h , and Δ^+ makes $d\Lambda$ dominant. Every relative discrete series representation is obtained this way, and $\pi(\gamma) \simeq \pi(\gamma')$ if and only if γ and γ' are K -conjugate.

The L-packet of γ is

$$(4.5) \quad \Pi_G(\gamma) = \{\pi(w\gamma) \mid w \in W(G, H)/W(K, H)\}.$$

It is a basic result of Harish-Chandra that $\pi(\gamma)$ exists and is unique. This version of the character formula is a slight variant of the usual one, because of the use of $\widetilde{H}(\mathbb{R})$. See [5] or [2].

By (4.4) the representations in $\Pi_G(\gamma)$ all have infinitesimal character $d\Lambda$. If $\rho = \rho_i$ is one-half the sum of any choice of positive roots (all roots are imaginary), $\Lambda \otimes e^\rho$ factors to $H(\mathbb{R})$, and the central character of $\Pi_G(\gamma)$ is $(\Lambda \otimes e^\rho)|_{Z(G(\mathbb{R}))}$.

Since $2\rho = 2\rho_i$ is a sum of roots, $e^{2\rho}$ is trivial on the center Z of G , and there is a canonical splitting of the restriction of \widetilde{H} to Z : $z \rightarrow (z, 1) \in H_\rho \simeq \widetilde{H}$. Using this splitting the central character of the packet is simply $\Lambda|_{Z(G(\mathbb{R}))}$.

Furthermore $\Pi(\gamma)$ is precisely the set of relative discrete series representations with the same infinitesimal and central characters as $\pi(\gamma)$. In fact something stronger is true.

Let G_{rad} be the radical of G . This is the maximal central torus, is the identity component of the center, and is defined over \mathbb{R} . By a *radical character* we mean a character of $G_{\text{rad}}(\mathbb{R})$, and the radical character of an irreducible representation is the restriction of its central character to $G_{\text{rad}}(\mathbb{R})$.

Proposition 4.6 *An L-packet of relative discrete series representations is uniquely determined by an infinitesimal and a radical character.*

This is based on the following structural fact.

Lemma 4.7 *Suppose $H(\mathbb{R})$ is a relatively compact Cartan subgroup of $G(\mathbb{R})$. Then*

$$(4.8) \quad Z(G(\mathbb{R})) \subset G_{\text{rad}}(\mathbb{R})H(\mathbb{R})^0.$$

Proof. Let $H_d = H \cap G_d$. Then $H_d(\mathbb{R})$ is a compact torus, and is therefore connected. It is enough to show:

$$(4.9) \quad H(\mathbb{R}) = G_{\text{rad}}(\mathbb{R})H_d(\mathbb{R}),$$

since this implies $Z(G(\mathbb{R})) \subset H(\mathbb{R}) = G_{\text{rad}}(\mathbb{R})H_d(\mathbb{R}) = G_{\text{rad}}(\mathbb{R})H(\mathbb{R})^0$. It is well known (4.9) holds over \mathbb{C} : $H(\mathbb{C}) = G_{\text{rad}}(\mathbb{C})H_d(\mathbb{C})$. If $y \in H(\mathbb{R})$ choose $z \in G_{\text{rad}}(\mathbb{C}), h \in H_d(\mathbb{C})$ such that $y = zh$. Since $y \in H(\mathbb{R}), \sigma(zh) = zh$, so

$$(4.10) \quad z\sigma(z^{-1}) = h^{-1}\sigma(h).$$

The left hand side is in $Z(G(\mathbb{C}))$, and the right hand side is in $G_d(\mathbb{C})$. So $h^{-1}\sigma(h) \in Z(G_d(\mathbb{C}))$.

Lemma 4.11 *Suppose $G(\mathbb{C})$ is a complex, connected, semisimple group, σ is an antiholomorphic involution of $G(\mathbb{C})$, $H(\mathbb{C})$ is a σ -stable Cartan subgroup, and $H(\mathbb{C})^\sigma = H(\mathbb{R})$ is compact. Then for $h \in H(\mathbb{C})$:*

$$(4.12) \quad h^{-1}\sigma(h) \in Z(G(\mathbb{C})) \Rightarrow h^{-1}\sigma(h) = 1$$

Proof. Let $G_{\text{ad}}(\mathbb{C})$ be the adjoint group and let $H_{\text{ad}}(\mathbb{C})$ be the image of $H(\mathbb{C})$ in $G_{\text{ad}}(\mathbb{C})$. Write p for projection $H(\mathbb{C}) \rightarrow H_{\text{ad}}(\mathbb{C})$. The condition $h^{-1}\sigma(h) \in Z(G(\mathbb{C}))$ is equivalent to $p(h) \in H_{\text{ad}}(\mathbb{R})$. So we need to show:

$$p(h) \in H_{\text{ad}}(\mathbb{R}) \Rightarrow h \in H(\mathbb{R}),$$

or equivalently, $p^{-1}(H_{\text{ad}}(\mathbb{R})) = H(\mathbb{R})$.

Since $H_{\text{ad}}(\mathbb{R})$ is a compact torus it is connected, so $p : H(\mathbb{R}) \rightarrow H_{\text{ad}}(\mathbb{R})$ is surjective. On the other hand, since $G(\mathbb{C})$ is semisimple every element of $Z(G(\mathbb{C}))$ has finite order. But $Z(G(\mathbb{C})) \subset H(\mathbb{C})$, and since $H(\mathbb{R})$ is compact this implies $Z(G(\mathbb{C})) \subset H(\mathbb{R})$. Therefore $p^{-1}(H_{\text{ad}}(\mathbb{R})) = H(\mathbb{R})Z(G(\mathbb{C})) = H(\mathbb{R})$. \square

By the Lemma applied to $G_d(\mathbb{C})$, $h^{-1}\sigma(h) = 1$, i.e. $h \in H_d(\mathbb{R})$, and $z \in G_{\text{rad}}(\mathbb{R})$ by (4.10). \square

Proof of the Proposition. Suppose Π is an L-packet of relative discrete series representations, with infinitesimal character χ_{inf} and central character χ . Choose $\lambda \in \mathfrak{h}^*$ defining χ_{inf} . It is enough to show χ is determined by λ and the restriction χ_{rad} of χ to $G_{\text{rad}}(\mathbb{R})$. But this is clear from Lemma 4.7: there is a unique character of $Z(G(\mathbb{R}))$ whose restriction to $G_{\text{rad}}(\mathbb{R})$ is χ_{rad} , and to $H(\mathbb{R})^0 \cap Z(G(\mathbb{R}))$ is $\exp(\lambda + \rho)$ (where ρ is one-half the sum of any set of positive roots). \square

The following converse to Proposition 4.6 follows immediately from the definitions.

Lemma 4.13 *Assume $G(\mathbb{R})$ has a relatively compact Cartan subgroup, and fix one, denoted $H(\mathbb{R})$. Suppose $\chi_{\text{inf}}, \chi_{\text{rad}}$ are infinitesimal and radical characters, respectively. Choose $\lambda \in \mathfrak{h}^*$ defining χ_{inf} via the Harish-Chandra homomorphism. Then the L-packet of relative discrete series representations defined by $\chi_{\text{inf}}, \chi_{\text{rad}}$ is nonzero if and only if λ is regular, and there is a genuine character of $\widetilde{H(\mathbb{R})}$ satisfying:*

$$(1) \quad d\Lambda = \lambda$$

$$(2) \quad \Lambda|_{G_{rad}(\mathbb{R})} = \chi_{rad}.$$

The conditions are independent of the choices of $H(\mathbb{R})$ and λ .

In (2) we have used the splitting $Z(G(\mathbb{R})) \rightarrow \widetilde{H(\mathbb{R})}$ discussed after (4.5).

We now describe general L-packets. See [2, Section 13] or [1, Section 6]

Definition 4.14 *Suppose γ is an L-datum. Let A be the identity component of $\{h \in H \mid \theta(h) = h^{-1}\}$ and set $M = \text{Cent}_G(A)$. Let $\mathfrak{a} = \text{Lie}(A)$. Choose a parabolic subgroup $P = MN$ satisfying*

$$\text{Re}\langle d\Lambda|_{\mathfrak{a}}, \vee\alpha \rangle \geq 0 \text{ for all roots of } \mathfrak{h} \text{ in } \text{Lie}(N).$$

Then P is defined over \mathbb{R} , and $H(\mathbb{R})$ is a relatively compact Cartan subgroup of $M(\mathbb{R})$. Let $\Pi_M(\gamma)$ be the L-packet of relative discrete series representations of $M(\mathbb{R})$ as in Definition 4.3. Define

$$(4.15) \quad \Pi_G(\gamma) = \bigcup_{\pi \in \Pi_M(\gamma)} \{\text{irreducible quotients of } \text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}(\pi)\}$$

Here we use normalized induction, and pull π back to $P(\mathbb{R})$ via the map $P(\mathbb{R}) \rightarrow M(\mathbb{R})$ as usual.

By the discussion following Definition 4.3, and basic properties of induction, the infinitesimal character of $\Pi_G(\gamma)$ is $d\Lambda$, and the central character is $\Lambda|_{Z(G(\mathbb{R}))}$.

5 The Tits Group

We need a few structural facts provided by the Tits group.

Fix a pinning $\mathcal{P} = (B, H, \{X_\alpha\})$ (see Section 2). For $\alpha \in \Pi$ define $X_{-\alpha} \in \mathfrak{g}_{-\alpha}$ by $[X_\alpha, X_{-\alpha}] = \vee\alpha$ as in Section 2. Define $\sigma_\alpha \in W = \exp(\frac{\pi}{2}(X_\alpha - X_{-\alpha})) \in \text{Norm}_G(H)$. The image of σ_α in W is the simple reflection s_α . Let $H_2 = \{h \in H \mid h^2 = 1\}$.

Definition 5.1 *The Tits group defined by \mathcal{P} is the subgroup \mathcal{T} of G generated by H_2 and $\{\sigma_\alpha \mid \alpha \in \Pi\}$.*

Proposition 5.2 ([13]) *The Tits group \mathcal{T} has the given generators, and relations:*

- (1) $\sigma_\alpha h \sigma_\alpha^{-1} = s_\alpha(h)$,
- (2) the braid relations among the σ_α ,
- (3) $\sigma_\alpha^2 = \vee_\alpha(-1)$.

If $w \in W$ then there is a canonical preimage σ_w of w in \mathcal{T} defined as follows. Suppose $w = s_{\alpha_1} \dots s_{\alpha_n}$ is a reduced expression with each $\alpha_i \in \Pi$. Then $\sigma_w = \sigma_{\alpha_1} \dots \sigma_{\alpha_n}$, independent of the choice of reduced expression.

Lemma 5.3 *If w_0 is the long element of the Weyl group, then σ_{w_0} is fixed by any \mathcal{P} -distinguished automorphism.*

Proof. Suppose $w_0 = s_{\alpha_1} \dots s_{\alpha_n}$ is a reduced expression. If γ is distinguished it induces an automorphism of the Dynkin diagram, so $s_{\gamma(\alpha_1)} \dots s_{\gamma(\alpha_n)}$ is also a reduced expression for w_0 . Therefore $\gamma(\sigma_{w_0}) = \gamma(\sigma_{\alpha_1} \dots \sigma_{\alpha_n}) = \sigma_{\gamma(\alpha_1)} \dots \sigma_{\gamma(\alpha_n)} = \sigma_{w_0}$ by the last assertion of Proposition 5.2. \square

Let \vee_ρ be one-half the sum of the positive coroots.

Lemma 5.4 *For any $w \in W$ we have*

$$(5.5) \quad \sigma_w \sigma_{w^{-1}} = \exp(\pi i(\vee_\rho - w \vee_\rho)).$$

In particular if w_0 is the long element of the Weyl group,

$$(5.6) \quad \sigma_{w_0}^2 = \exp(2\pi i \vee_\rho) \in Z(G).$$

This element of $Z(G)$ is independent of the choice of positive roots, and is fixed by every automorphism of G .

Proof. We proceed by induction on the length of w . If w is a simple reflection s_α then $s_\alpha \vee_\rho = \vee_\rho - \vee_\alpha$, and this reduces to Proposition 5.2(3).

Write $w = s_\alpha u$ with α simple and $\ell(w) = \ell(u) + 1$. Then $\sigma_w = \sigma_\alpha \sigma_u$, $w^{-1} = u^{-1} s_\alpha$, $\sigma_w = \sigma_{u^{-1}} \sigma_\alpha$, and

$$(5.7) \quad \begin{aligned} \sigma_w \sigma_{w^{-1}} &= \sigma_\alpha \sigma_u \sigma_{u^{-1}} \sigma_\alpha \\ &= \sigma_\alpha \sigma_u \sigma_{u^{-1}} \sigma_\alpha^{-1} m_\alpha \\ &= \sigma_\alpha \exp(\vee_\rho - u \vee_\rho) \sigma_\alpha^{-1} m_\alpha \quad (\text{by the inductive step}) \\ &= \exp(\pi i(s_\alpha \vee_\rho - w \vee_\rho)) \exp(\pi i \vee_\alpha) \\ &= \exp(\pi i(\vee_\rho - \vee_\alpha - w \vee_\rho)) \exp(\pi i \vee_\alpha) = \exp(\pi i(\vee_\rho - w \vee_\rho)). \end{aligned}$$

The final assertion is easy. \square

We thank Marc van Leeuwen for this proof.

We only need what follows for the part of the main theorem involving the Chevalley involution τ of $W_{\mathbb{R}}$. Let $C = C_{\mathcal{P}}$ be the Chevalley involution defined by \mathcal{P} .

Lemma 5.8 $C(\sigma_w) = (\sigma_{w^{-1}})^{-1}$

Proof. We proceed by induction on the length of w .

Since $C(X_{\alpha}) = X_{-\alpha}$ ($\alpha \in \Pi$), we conclude $C(\sigma_{\alpha}) = \sigma_{\alpha}^{-1}$.

Suppose $w = s_{\alpha}u$ with $\text{length}(w) = \text{length}(u) + 1$. Then $\sigma_w = \sigma_{\alpha}\sigma_u$, and $C(\sigma_w) = C(\sigma_{\alpha})C(\sigma_u) = \sigma_{\alpha}^{-1}(\sigma_{u^{-1}})^{-1}$. On the other hand $w^{-1} = u^{-1}s_{\alpha}$, so $\sigma_{w^{-1}} = \sigma_{u^{-1}}\sigma_{\alpha}$, and taking the inverse gives the result. \square

Fix a \mathcal{P} -distinguished involution τ of G . Consider the semidirect product $G \rtimes \langle \delta \rangle$ where δ acts on G by τ . By Lemma 2.6, $C = C_{\mathcal{P}}$ extends to the semidirect product, fixing δ . Since τ normalizes H , it defines an automorphism of W , satisfying $\tau(\sigma_w) = \sigma_{\tau(w)}$.

Lemma 5.9 *Suppose $w\tau(w) = 1$. Then*

(a) $C(\sigma_w\delta) = (\sigma_w\delta)^{-1}$

(b) *Suppose $g \in \text{Norm}_G(H)$ is a representative of w . Then $C(g\delta)$ is H -conjugate to $(g\delta)^{-1}$.*

Proof. By the previous Lemma, and using $\tau(w) = w^{-1}$, we compute

$$C(\sigma_w\delta)\sigma_w\delta = (\sigma_{w^{-1}})^{-1}\sigma_{\tau(w)} = (\sigma_{\tau(w)})^{-1}\sigma_{\tau(w)} = 1.$$

For (b) write $g = h\sigma_w\delta$ with $h \in H$. Then $C(g) = C(h\sigma_w\delta) = h^{-1}(\sigma_w\delta)^{-1} = h^{-1}(h\sigma_w\delta)^{-1}h$. \square

6 L-parameters

Fix a Cartan involution θ of G . Let ${}^{\vee}G$ be the connected, complex dual group of G . The L-group ${}^L G$ of G is $\langle {}^{\vee}G, {}^{\vee}\delta \rangle$ where ${}^{\vee}\delta^2 = 1$, and ${}^{\vee}\delta$ acts on ${}^{\vee}G$ by a homomorphism ${}^{\vee}\theta_0$, which we now describe. See [6], [5], or [4, Section 2].

Fix B_0, H_0 and let $D_b = (X^*(H_0), \Pi, X_*(H_0), \vee\Pi)$ be the corresponding based root datum. Similarly choose $\vee B_0, \vee H_0$ for $\vee G$ to define $\vee D_b$. In particular we identify $X^*(H_0) = X_*(\vee H_0)$ and $X_*(H_0) = X^*(\vee H_0)$. Also fix a pinning $\vee\mathcal{P} = (\vee B_0, \vee H_0, \{X_{\vee\alpha}\})$ for $\vee G$. See Section 2.

An automorphism γ of D_b consists of a pair (τ, τ^t) , where $\tau \in \text{Aut}(X^*(H_0))$, and $\tau^t \in \text{Aut}(X_*(H_0))$ is the transpose with respect to the perfect pairing $X^*(H_0) \times X_*(H_0) \rightarrow \mathbb{Z}$. These preserve $\Pi, \vee\Pi$ respectively. Interchanging (τ, τ^t) defines a transpose isomorphism $\text{Aut}(D_b) \simeq \text{Aut}(\vee D_b)$, denoted $\mu \rightarrow \mu^t$. Compose with the embedding $\text{Aut}(\vee D_b) \hookrightarrow \text{Aut}(\vee G)$ defined by $\vee\mathcal{P}$ (Section 2) to define a map:

$$(6.1) \quad \mu \rightarrow \mu^t : \text{Aut}(D_b) \simeq \text{Aut}(\vee D_b) \hookrightarrow \text{Aut}(\vee G).$$

Here are two equivalent ways of defining ${}^L G$, each of which has its advantages.

Let γ be the image of θ in $\text{Out}(G) \simeq \text{Aut}(D_b)$. The long element w_0 of $W(G, H_0)$ satisfies $-w_0 \in \text{Aut}(D_b)$. Define

$$(6.2) \quad \vee\theta_0 = (-w_0\gamma)^t \in \text{Aut}(\vee G).$$

This is the approach of [5] and [4].

Alternatively suppose σ is a real form corresponding to θ (see the beginning of Section 4). Choose g conjugating $\sigma(B_0)$ to B_0 and $\sigma(H_0)$ to H_0 . Then $\tau = \text{int}(g) \circ \sigma \in \text{Aut}(D_b)$. Let $\vee\theta_0 = \tau^t \in \text{Aut}(\vee G)$. See [6].

It is not hard to see the elements $\tau, \gamma \in \text{Aut}(D_b)$ satisfy $\tau = -w_0\gamma$, so the two definitions of ${}^L G$ agree.

The second definition is well suited to the most split Cartan subgroup. For example suppose $G(\mathbb{R})$ is split. Choosing B_0, H_0 defined over \mathbb{R} gives $\tau = 1$, so $\tau^t = 1$ and ${}^L G = \vee G \times \Gamma$ (direct product).

On the other hand the first version is natural with respect to the fundamental (most compact) Cartan subgroup. The next Lemma is very useful in this regard.

Lemma 6.3 *The following conditions are equivalent:*

- (1) $G(\mathbb{R})$ has a compact Cartan subgroup,
- (2) $\vee\theta_0$ is inner to the Chevalley involution;

(3) There is an element $y \in {}^L G \backslash {}^\vee G$ such that $yhy^{-1} = h^{-1}$ for all $y \in {}^\vee H_0$.

Proof. The equivalence of (2) and (3) is immediate. Let C be the Chevalley involution of ${}^\vee G$ with respect to ${}^\vee \mathcal{P}$. It is easy to see (6.2) is equivalent to: the image of ${}^\vee \theta_0 \circ C$ in $\text{Out}({}^\vee G) \simeq \text{Aut}({}^\vee D_b)$ is equal to γ^t . So the assertion is that $G(\mathbb{R})$ has a compact Cartan subgroup if and only if $\gamma = 1$, i.e. θ is an inner automorphism, which holds by Lemma 6.3. \square

A homomorphism $\phi : W_{\mathbb{R}} \rightarrow {}^L G$ is said to be *admissible* if it is continuous, $\phi(\mathbb{C}^*)$ consists of semisimple elements, and $\phi(j) \in {}^L G \backslash {}^\vee G$ [6, 8.2]. Associated to an admissible homomorphism ϕ is an L-packet $\Pi_G(\phi)$, which depends only on the ${}^\vee G$ -conjugacy class of ϕ . Since we did not impose the relevancy condition [6, 8.2(ii)], such a packet may be empty.

After conjugating by ${}^\vee G$ we may assume $\phi(\mathbb{C}^*) \subset {}^\vee H_0$. Let ${}^\vee S$ be the centralizer of $\phi(\mathbb{C}^*)$ in ${}^\vee G$. Since $\phi(\mathbb{C}^*)$ is connected, abelian and consists of semisimple elements, ${}^\vee S$ is a connected reductive complex group, and ${}^\vee H_0$ is a Cartan subgroup of ${}^\vee S$. Conjugation by $\phi(j)$ is an involution of ${}^\vee S$, so $\phi(j)$ normalizes a Cartan subgroup of ${}^\vee S$. Equivalently some ${}^\vee S$ -conjugate of $\phi(j)$ normalizes ${}^\vee H_0$; after this change we may assume $\phi(W_{\mathbb{R}}) \subset \text{Norm}_{{}^\vee G}({}^\vee H_0)$.

Therefore

$$(6.4)(a) \quad \phi(z) = z^\lambda \bar{z}^{\lambda'} \quad (\text{for some } \lambda, \lambda' \in X_*({}^\vee H_0) \otimes \mathbb{C}, \lambda - \lambda' \in X_*({}^\vee H_0))$$

$$(6.4)(b) \quad \phi(j) = h\sigma_w{}^\vee \delta \quad (\text{for some } w \in W, h \in {}^\vee H_0).$$

Here (a) is shorthand for $\phi(e^s) = \exp(s\lambda + \bar{s}\lambda') \in {}^\vee H_0$ ($s \in \mathbb{C}$), and the condition on $\lambda - \lambda'$ guarantees this is well defined. In (b) we're using the element σ_w of the Tits group representing w (Proposition 5.2).

Conversely, given λ, λ', w and h , (a) and (b) give a well-defined homomorphism $\phi : W_{\mathbb{R}} \rightarrow {}^L G$ if and only if

$$(6.4)(c) \quad {}^\vee \theta := \text{int}(h\sigma_w{}^\vee \delta) \text{ is an involution of } {}^\vee H_0,$$

$$(6.4)(d) \quad \lambda' = {}^\vee \theta(\lambda),$$

$$(6.4)(e) \quad h{}^\vee \theta(h)(\sigma_w{}^\vee \theta_0(\sigma_w)) = \exp(\pi i(\lambda - {}^\vee \theta(\lambda))).$$

Furthermore (c) is equivalent to

$$(6.4)(c') \quad w{}^\vee \delta(w) = 1.$$

6.1 Infinitesimal and radical characters

Infinitesimal Character of $\Pi_G(\phi)$

View λ as an element of $X^*(H_0) \otimes \mathbb{C}$ via the identification $X_*(\vee H_0) = X^*(H_0)$. The $W(G, H)$ -orbit of λ is independent of all choices, so it defines an infinitesimal character for G , denoted $\chi_{\text{inf}}(\phi)$, via the Harish-Chandra homomorphism.

Radical character of $\Pi(\phi)$

Recall (Section 4) G_{rad} is the radical of G , and the radical character of a representation is its restriction to $G_{\text{rad}}(\mathbb{R})$.

Dual to the inclusion $\iota : G_{\text{rad}} \hookrightarrow G$ is a surjection $\vee \iota : \vee G \rightarrow \vee[G_{\text{rad}}]$ (the dual group of G_{rad}). For an L-group for G_{rad} we can take ${}^L G_{\text{rad}} = \langle \vee G_{\text{rad}}, \vee \delta \rangle$. Thus $\vee \iota$ extends to a natural surjection $\vee \iota : {}^L G \rightarrow {}^L G_{\text{rad}}$ (taking $\vee \delta$ to itself). Then $\vee \iota \circ \phi : W_{\mathbb{R}} \rightarrow {}^L G_{\text{rad}}$, and this defines a character of $G_{\text{rad}}(\mathbb{R})$ by the construction of Section 3. We denote this character $\chi_{\text{rad}}(\phi)$. See [6, 10.1]

6.2 Relative Discrete Series L-packets

By a *Levi subgroup* of ${}^L G$ we mean the centralizer ${}^d M$ of a torus $\vee T \subset \vee G$, which meets both components of ${}^L G$ [6, Lemma 3.5]. An L-packet $\Pi_G(\phi)$ consists of relative discrete series representations if and only if $\phi(W_{\mathbb{R}})$ is not contained in a proper Levi subgroup.

Lemma 6.5 *$\phi(W_{\mathbb{R}})$ is not contained in a proper Levi subgroup if and only if λ is regular, and $G(\mathbb{R})$ has a relatively compact Cartan subgroup.*

Proof. Assume $\phi(W_{\mathbb{R}})$ is not contained in a proper Levi subgroup. Let $\vee S = \text{Cent}_{\vee G}(\phi(W_{\mathbb{R}}))$ as in the discussion preceding (6.4). Then $\vee \theta = \text{int}(\phi(j))$ is an involution of $\vee S$, and of its derived group $\vee S_d$. There cannot be a torus in $\vee S_d$, fixed (pointwise) by $\vee \theta$; its centralizer would contradict the assumption. Since any involution of a semisimple group fixes a torus, this implies $\vee S_d = 1$, i.e. $\vee S = \vee H_0$, which implies λ is regular.

Similarly, there can be no torus in $\vee H_0 \cap \vee G_d$ fixed by $\vee \theta$. This implies $\vee \theta(h) = h^{-1}$ for all $h \in \vee H_0 \cap \vee G_d$. By Lemma 6.3 applied to the derived group, $G(\mathbb{R})$ has a relatively compact Cartan subgroup.

The reverse implication is similar. □

Definition 6.6 $\Pi_G(\phi)$ is the L -packet of relative discrete series representations determined by infinitesimal character $\chi_{\text{inf}}(\phi)$ and radical character $\chi_{\text{rad}}(\phi)$ (see Proposition 4.6).

Lemma 6.7 $\Pi(\phi)$ is nonempty.

Proof. After conjugating θ if necessary we may assume H_0 is θ -stable. Write ϕ as in (6.4)(a) and (b). By the discussion of infinitesimal character above $\chi_{\text{inf}}(\phi)$ is defined by λ , viewed as an element of $X^*(H_0) \otimes \mathbb{C}$.

Choose positive roots making λ dominant. By Lemma 4.13 it is enough to construct a genuine character of $H(\mathbb{R})_\rho$ satisfying $d\Lambda = \lambda$ and $\Lambda|_{G_{\text{rad}}(\mathbb{R})} = \chi_{\text{rad}}$. For this we apply Lemma 3.3, using the fact that $\phi(W_{\mathbb{R}}) \subset \langle {}^\vee H_0, \sigma_w {}^\vee \delta \rangle$. First we need to identify $\langle {}^\vee H_0, \sigma_w {}^\vee \delta \rangle$ as an E-group.

First we claim $w = w_0$. By (6.4)(b) and (c) ${}^\vee \theta|_{{}^\vee H_0} = w {}^\vee \theta_0|_{{}^\vee H_0}$. By (6.2), for $h \in {}^\vee H_0 \cap {}^\vee G_d$ we have:

$$(6.8) \quad {}^\vee \theta(h) = w {}^\vee \theta_0(h) = w(-w_0 \gamma^t)(h) = w w_0 \gamma^t(h)^{-1}.$$

Since $G_d(\mathbb{R})$ has a compact Cartan γ is trivial on $H_0 \cap G_d$ and $\gamma^t(h) = h$. On the other hand, as in the proof of Lemma 6.5, ${}^\vee \theta(h) = h^{-1}$. Therefore $w w_0 = 1$, i.e. $w = w_0$.

Next we compute

$$(6.9) \quad \begin{aligned} (\sigma_{w_0} {}^\vee \delta)^2 &= w_0 \theta_0(w_0) \\ &= w_0^2 \quad (\text{by (5.3), since } {}^\vee \theta_0 \text{ is distinguished}) \\ &= \exp(2\pi i \rho) \quad (\text{by (5.6)}). \end{aligned}$$

Remark 6.10 The fact that $(\sigma_{w_0} {}^\vee \delta)^2 = \exp(2\pi i \rho)$ is the analogue of [10, Lemma 3.2].

Thus, in the terminology of Section 3, $\langle {}^\vee H_0, \sigma_{w_0} {}^\vee \delta \rangle$ is identified with the E-group for H defined by ${}^\vee \rho$. Thus $\phi : W_{\mathbb{R}} \rightarrow \langle {}^\vee H_0, \sigma_{w_0} {}^\vee \delta \rangle$ defines a genuine character Λ of $H(\mathbb{R})_\rho$.

By construction $d\Lambda = \lambda$. The fact that $\Lambda|_{G_{\text{rad}}(\mathbb{R})} = \chi_{\text{rad}}$ is a straightforward check. Here are the details.

Write h of (6.4)(b) as $h = \exp(2\pi i \mu)$. We use the notation of Section 3, especially (3.4). Using the fact that $\sigma_{w_0} {}^\vee \delta$ is the distinguished element of the E-group of H_0 we have $\Lambda = \chi(\lambda, \kappa)$ where

$$\kappa = \frac{1}{2}(1 - {}^\vee \theta)\lambda - (1 + {}^\vee \theta)\mu \in \rho + X^*(H_0).$$

Write $p : X^*(H_0) \rightarrow X^*(G_{\text{rad}})$ for the map dual to inclusion $G_{\text{rad}} \rightarrow H_0$. Recall (Section 4) there is a canonical splitting of the cover of $H(\mathbb{R})$; using this splitting $\chi(\rho, \rho)|_{Z(G(\mathbb{R}))} = 1$, and

$$\Lambda|_{G_{\text{rad}}(\mathbb{R})} = \chi(p(\lambda), p(\kappa - \rho)).$$

On the other hand, by the discussion of the character of $G_{\text{rad}}(\mathbb{R})$ above, the E-group of G_{rad} is $\langle {}^\vee G_{\text{rad}}, {}^\vee \delta \rangle$. The map $p : X^*(H_0) \rightarrow X^*(G_{\text{rad}})$ is identified with a map $p : X_*({}^\vee H_0) \rightarrow X_*({}^\vee G_{\text{rad}})$. Then ${}^\vee \iota(\phi(j)) = {}^\vee \iota(h\sigma_w {}^\vee \delta) = {}^\vee \iota(h) {}^\vee \delta = \exp(2\pi i p(\mu)) {}^\vee \delta$. Let

$$\kappa' = \frac{1}{2}(1 - {}^\vee \theta)p(\lambda) - (1 + {}^\vee \theta)p(\mu) \in X^*(G_{\text{rad}})$$

Thus $\kappa' = p(\kappa - \rho)$. By the construction of Section 3 applied to $\phi : W_{\mathbb{R}} \rightarrow {}^\vee G_{\text{rad}}$, $\chi_{\text{rad}} = \chi(p(\lambda), \kappa') = \chi(p(\lambda), p(\kappa - \rho)) = \Lambda|_{G_{\text{rad}}(\mathbb{R})}$. \square

We can read off the central character of the L-packet from the construction. We defer this until we consider general L-packets (Lemma 6.16).

6.3 General L-packets

Recall (see the beginning of the previous section) a Levi subgroup ${}^d M$ of ${}^L G$ is the centralizer of a torus ${}^\vee T$, which meets both components of ${}^L G$. An admissible homomorphism ϕ may factor through various Levi subgroups ${}^d M$. We first choose ${}^d M$ so that $\phi : W_{\mathbb{R}} \rightarrow {}^L M$ defines a relative discrete series L-packet of M .

Choose a maximal torus ${}^\vee T \subset \text{Cent}_{{}^\vee G}(\phi(W_{\mathbb{R}}))$ and define

$$(6.11) \quad {}^\vee M = \text{Cent}_{{}^\vee G}({}^\vee T), \quad {}^d M = \text{Cent}_{{}^L G}({}^\vee T).$$

Then ${}^d M = \langle {}^\vee M, \phi(j) \rangle$, so ${}^d M$ is a Levi subgroup, and $\phi(W_{\mathbb{R}}) \subset {}^d M$.

Suppose $\phi(W_{\mathbb{R}}) \subset \text{Cent}_{{}^d M}({}^\vee U)$ where ${}^\vee U \subset {}^\vee M$ is a torus. Then ${}^\vee U$ centralizes $\phi(W_{\mathbb{R}})$ and ${}^\vee T$, so ${}^\vee U {}^\vee T$ is a torus in $\text{Cent}_{{}^\vee G}(\phi(W_{\mathbb{R}}))$. By maximality ${}^\vee U \subset {}^\vee T$ and $\text{Cent}_{{}^d M}({}^\vee U) = {}^d M$. Therefore $\phi(W_{\mathbb{R}})$ is not contained in any proper Levi subgroup of ${}^d M$.

Lemma 6.12 *The group dM is independent of the choice of ${}^\vee T$, up to conjugation by $\text{Cent}_{\vee G}(\phi(W_{\mathbb{R}}))$.*

Proof. Let ${}^\vee C = \text{Cent}_{\vee G}(\phi(W_{\mathbb{R}}))$. Conjugation by $\phi(j)$ is an involution of the connected reductive group ${}^\vee S = \text{Cent}_{\vee G}(\phi(\mathbb{C}^*))$ (see the discussion after Lemma 6.3) and ${}^\vee C$ is the fixed points of this involution. Therefore ${}^\vee C$ is a reductive group, and ${}^\vee T$ is a maximal torus in the identity component ${}^\vee C^0$. Any two such tori are conjugate by ${}^\vee C^0$. \square

The idea is to identify dM with the L-group of a Levi subgroup $M'(\mathbb{R})$ of $G(\mathbb{R})$. Then, since $\phi(W_{\mathbb{R}}) \subset {}^dM$ is not contained in any proper Levi subgroup, it defines a relative discrete series L-packet for $M'(\mathbb{R})$. We obtain Π_G by induction. Here are the details.

We need to identify ${}^dM = \langle {}^\vee M, \phi(j) \rangle$ with an L-group. A crucial technical point is that after conjugating we may assume ${}^dM = \langle {}^\vee M, {}^\vee \delta \rangle$, making this identification clear.

Lemma 6.13 ([6], Section 3.1) *Suppose S is a ${}^\vee\theta_0$ -stable subset of ${}^\vee\Pi$. Let ${}^\vee M_S$ be the corresponding Levi subgroup of ${}^\vee G$: ${}^\vee H_0 \subset {}^\vee M_S$, and S is a set of simple roots of ${}^\vee H_0$ in ${}^\vee M_S$. Let ${}^L M_S = {}^\vee M_S \rtimes \langle {}^\vee \delta \rangle$, a Levi subgroup of ${}^L G$.*

Let $M_S \supset H_0$ be the Levi subgroup of G with simple roots $\{\alpha \mid {}^\vee \alpha \in S\} \subset \Pi$. Given a real form of G (in the given inner class) suppose some conjugate M' of M_S is defined over \mathbb{R} . Write ${}^L M' = {}^\vee M' \rtimes \langle {}^\vee \delta_{M'} \rangle$. Then conjugation induces an isomorphism ${}^L M' \simeq {}^L M_S$, taking ${}^\vee \delta_{M'}$ to ${}^\vee \delta$.

Any Levi subgroup of ${}^L G$ is ${}^\vee G$ -conjugate to ${}^L M_S$ for some ${}^\vee\theta_0$ -stable set S .

We refer to the Levi subgroups ${}^L M_S$ of the Lemma (where S is ${}^\vee\theta_0$ -stable) as standard Levi subgroups.

Definition 6.14 *Suppose $\phi : W_{\mathbb{R}} \rightarrow {}^L G$ is an admissible homomorphism. Choose a maximal torus ${}^\vee T$ in $\text{Cent}_{\vee G}(\phi(W_{\mathbb{R}}))$, and define ${}^\vee M, {}^dM$ by (6.11).*

After conjugating by ${}^\vee G$, we we may assume dM is a standard Levi subgroup. Let M be the corresponding standard Levi subgroup of G .

Assume there is a subgroup M' conjugate to M , which is defined over \mathbb{R} ; otherwise $\Pi(\phi)$ is empty. Let $\Pi_{M'}(\phi)$ be the L-packet for $M'(\mathbb{R})$ defined by

$\phi : W_{\mathbb{R}} \rightarrow {}^d M \simeq {}^L M'$. (cf. Lemma 6.13). Define $\Pi_G(\phi)$ by induction from $\Pi_{M'}(\phi)$ as in Definition 4.14.

Lemma 6.15 *The L-packet $\Pi_G(\phi)$ is independent of all choices.*

Proof. By Lemma 6.12 the choice of ${}^V T$ is irrelevant: another choice leads to an automorphism of ${}^d M$ fixing $\phi(W_{\mathbb{R}})$ pointwise.

It is straightforward to see that the other choices, including another Levi subgroup M'' , would give an element $g \in G(\mathbb{C})$ such that $\text{int}(g) : M' \rightarrow M''$ is defined over \mathbb{R} , and this isomorphism takes $\Pi_{M'}(\phi)$ to $\Pi_{M''}(\phi)$.

First of all we claim $M'(\mathbb{R})$ and $M''(\mathbb{R})$ are $G(\mathbb{R})$ -conjugate. To see this, let $H'(\mathbb{R})$ be a relatively compact Cartan subgroup of $M'(\mathbb{R})$. Then $H''(\mathbb{R}) = gH'(\mathbb{R})g^{-1}$ is a relatively compact Cartan subgroup of $M''(\mathbb{R})$. In fact $H'(\mathbb{R})$ and $H''(\mathbb{R})$ are $G(\mathbb{R})$ -conjugate: two Cartan subgroups of $G(\mathbb{R})$ are $G(\mathbb{C})$ -conjugate if and only if they are $G(\mathbb{R})$ -conjugate. Therefore $M'(\mathbb{R}), M''(\mathbb{R})$, being the centralizers of the split components of $H'(\mathbb{R})$ and $H''(\mathbb{R})$, are also $G(\mathbb{R})$ -conjugate.

Therefore, since the inductive step is not affected by conjugating by $G(\mathbb{R})$, we may assume $M' = M''$. Then $g \in \text{Norm}_{G(\mathbb{C})}(M')$, and furthermore $g \in \text{Norm}_{G(\mathbb{C})}(M'(\mathbb{R}))$.

Now $gH'(\mathbb{R})g^{-1}$ is another relatively compact Cartan subgroup of $M'(\mathbb{R})$, so after replacing g with gm for some $m \in M'(\mathbb{R})$ we may assume $g \in \text{Norm}_{G(\mathbb{C})}(H'(\mathbb{R}))$. It is well known that

$$\text{Norm}_{G(\mathbb{C})}(H'(\mathbb{R})) = \text{Norm}_{M'(\mathbb{C})}(H'(\mathbb{R}))\text{Norm}_{G(\mathbb{R})}(H'(\mathbb{R})).$$

For example see [14, Proposition 3.12] (where the group in question is denoted $W(R)^\theta$), or [11, Theorem 2.1]. Since conjugation by $M'(\mathbb{C})$ does not change infinitesimal or central characters, by Proposition 4.6 it preserves $\Pi_{M'}(\phi)$. As above $G(\mathbb{R})$ has no effect after the inductive step. This completes the proof. \square

We now give the formula for the central character of $\Pi_G(\phi)$.

Lemma 6.16 *Write ϕ as in (6.4)(a) and (b), and suppose $h = \exp(2\pi i\mu)$, with $\mu \in X_*({}^V H_0) \otimes \mathbb{C} \simeq X^*(H_0) \otimes \mathbb{C}$. Let ρ_i be one-half the sum of any set of positive roots of $\{\alpha \mid {}^V \theta \alpha = -\alpha\}$. Set*

$$\tau = \frac{1}{2}(1 - {}^V \theta)\lambda - (1 + {}^V \theta)\mu + \rho_i \in X^*(H_0).$$

Then the central character of $\Pi_G(\phi)$ is $\tau|_{Z(G(\mathbb{R}))}$.

7 Contragredient

The proof of Theorem 1.3 is now straightforward. We restate the Theorem here.

Theorem 7.1 *Let $G(\mathbb{R})$ be the real points of a connected reductive algebraic group defined over \mathbb{R} , with L -group ${}^L G$. Let C be the Chevalley involution of ${}^L G$ (Section 2) and let τ be the Chevalley involution of $W_{\mathbb{R}}$ (Lemma 2.7). Suppose $\phi : W_{\mathbb{R}} \rightarrow {}^L G$ is an admissible homomorphism, with associated L -packet $\Pi(\phi)$. Let $\Pi(\phi)^* = \{\pi^* \mid \pi \in \Pi(\phi)\}$. Then:*

(a) $\Pi(\phi)^* = \Pi(C \circ \phi)$

(b) $\Pi(\phi)^* = \Pi(\phi \circ \tau)$

Proof. Let $\mathcal{P} = (\vee B_0, \vee H_0, \{X_{\vee\alpha}\})$ be the pinning used to define ${}^L G$. After conjugating by $\vee G$ we may assume $C = C_{\mathcal{P}}$. As in the discussion before (6.4), we are free to conjugate ϕ so that $\phi(\mathbb{C}^*) \in \vee H_0$, and $\phi(W_{\mathbb{R}}) \subset \text{Norm}_{\vee G}(\vee H_0)$.

First assume $\Pi(\phi)$ is an L -packet of relative discrete series representations. Then $\Pi(\phi)$ is determined by its infinitesimal character $\chi_{\text{inf}}(\phi)$ and its radical character $\chi_{\text{rad}}(\phi)$ (Definition 6.6). It is easy to see that the infinitesimal character of $\Pi(\phi)^*$ is $-\chi_{\text{inf}}(\phi)$, and the radical character is $\chi_{\text{rad}}(\phi)^*$. So it is enough to show $\chi_{\text{inf}}(C \circ \phi) = -\chi_{\text{inf}}(\phi)$ and $\chi_{\text{rad}}(C \circ \phi) = \chi_{\text{rad}}(\phi)^*$. The first is obvious from (6.4)(a), the Definition of $\chi_{\text{inf}}(\phi)$, and the fact that C acts by -1 on the Lie algebra of $\vee H_0$. The second follows from the fact that C factors to the Chevalley involution of ${}^L G_{\text{rad}}$, and the torus case (Lemma 3.6).

Now suppose ϕ is any admissible homomorphism such that $\Pi_G(\phi)$ is nonempty. As in Definition 4.14 we may assume $\phi(W_{\mathbb{R}}) \subset {}^d M$ where ${}^d M$ is a standard Levi subgroup of ${}^L G$. Choose M' as in Definition 6.14 and write $\Pi_{M'} = \Pi_{M'}(\phi)$ as in that Definition.

Write socle (resp. co-socle) for the set of irreducible submodules (resp. quotients) of an admissible representation.

Choose $P = M'N$ as in Definition 4.14 to define

$$(7.2)(a) \quad \Pi_G(\phi) = \text{cosocle}(\text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}(\Pi_{M'})) = \bigcup_{\pi \in \Pi_{M'}} \text{cosocle}(\text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}(\pi)).$$

It is immediate from the definitions that $C_{\mathcal{P}}$ restricts to the Chevalley involution of $\vee M$. Therefore by the preceding case $\Pi_{M'}(C \circ \phi) = \Pi_{M'}(\phi)^*$.

Apply Definition 4.14 again to compute $\Pi_G(C \circ \phi)$; this time the positivity condition in Definition 4.14 forces us to use the opposite parabolic $\bar{P} = M'\bar{N}$:

$$(7.2)(b) \quad \Pi_G(C \circ \phi) = \text{cosocle}(\text{Ind}_{\bar{P}(\mathbb{R})}^{G(\mathbb{R})}(\Pi_{M'}^*)).$$

Here is the proof of part (a), we justify the steps below.

$$(7.2)(c) \quad \begin{aligned} \Pi_G(C \circ \phi)^* &= [\text{cosocle}(\text{Ind}_{\bar{P}(\mathbb{R})}^{G(\mathbb{R})}(\Pi_{M'}^*))]^* \\ &= [\text{cosocle}(\text{Ind}_{\bar{P}(\mathbb{R})}^{G(\mathbb{R})}(\Pi_{M'})^*)]^* \\ &= [\text{socle}(\text{Ind}_{\bar{P}(\mathbb{R})}^{G(\mathbb{R})}(\Pi_{M'}))^*]^* \\ &= \text{cosocle}(\text{Ind}_{\bar{P}(\mathbb{R})}^{G(\mathbb{R})}(\Pi_{M'})) = \Pi_G(\phi). \end{aligned}$$

The first step is just the contragredient of (7.2)(b). For the second, integration along $G(\mathbb{R})/P(\mathbb{R})$ defines a pairing between $\text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}(\pi^*)$ and $\text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}(\pi)^*$, and gives

$$(7.3) \quad \text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}(\pi^*) \simeq \text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}(\pi)^*.$$

For the next step use $\text{cosocle}(X^*) = \text{socle}(X)^*$. Then the double dual cancels for irreducible representations, and it is well known that the theory of intertwining operators gives:

$$(7.4) \quad \text{socle}(\text{Ind}_{\bar{P}(\mathbb{R})}^{G(\mathbb{R})}(\pi)) = \text{cosocle}(\text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}(\pi)).$$

Finally plugging in (7.2)(a) gives part (a) of the Theorem.

For (b) we show that $C \circ \phi$ is ${}^V G$ -conjugate to $\phi \circ \tau$.

Recall τ is any automorphism of $W_{\mathbb{R}}$ acting by inverse on \mathbb{C}^* , and any two such τ are conjugate by $\text{int}(z)$ for $z \in \mathbb{C}^*$. Therefore the statement is independent of the choice of τ . It is convenient to choose $\tau(j) = j^{-1}$, i.e. $\tau = \tau_{-1}$ in the notation of the proof of Lemma 2.7.

By (6.4)(a) $(C \circ \phi)(z) = C(\phi(z)) = z^{-\lambda} \bar{z}^{-\lambda'}$. On the other hand $(\phi \circ \tau)(z) = \phi(z^{-1}) = z^{-\lambda} \bar{z}^{-\lambda'}$. Therefore it is enough to show $C(\phi(j))$ is ${}^V H_0$ -conjugate to $\phi(\tau(j))$, which equals $\phi(j)^{-1}$ by our choice of τ .

Since $\phi(j)$ normalizes ${}^V H_0$, $\phi(j) = g {}^V \delta$ with $g \in \text{Norm}_G({}^V H_0)$. Then $g \delta(g) \phi(j)^2 = \phi(-1) \in {}^V H_0$. Therefore the image w of g in w satisfies $w \theta_0(w) = 1$. Apply Lemma 5.9(b). \square

8 Hermitian Dual

Suppose π is an admissible representation of $G(\mathbb{R})$. We briefly recall what it means for π to have an invariant Hermitian form, and the notion of the Hermitian dual of π . See [9] for details, and for the connection with unitary representations.

We work entirely in a purely algebraic setting. Write σ for complex conjugation of $\mathfrak{g} = \text{Lie}(G(\mathbb{C}))$ with respect to $\mathfrak{g}_0 = \text{Lie}(G(\mathbb{R}))$. Choose a Cartan involution θ commuting with σ , and let $K = G^\theta$. We consider (\mathfrak{g}, K) -modules, and write (π, V) for a (\mathfrak{g}, K) -module with underlying complex vector space V .

We say (π, V) , or simply π , is Hermitian if there is a nondegenerate Hermitian form (\cdot, \cdot) on V , satisfying

$$(8.1) \quad (\pi(X)v, w) + (v, \pi(\sigma(X))w) = 0 \quad (v, w \in V, X \in \mathfrak{g}),$$

and a similar identity for the action of K .

Define the Hermitian dual (π^h, V^h) as follows. Define a representation of \mathfrak{g} on the space of conjugate-linear functions $V \rightarrow \mathbb{C}$ by

$$(8.2) \quad \pi^h(X)(f)(v) = -f(\pi(\sigma(X))v) \quad (v \in V, X \in \mathfrak{g}).$$

Define the action of K by a similar formula, and let V^h be the K -finite functions; then (π^h, V^h) is a (\mathfrak{g}, K) -module. If π is irreducible then π is Hermitian if and only if $\pi \simeq \pi^h$.

Fix a Cartan subgroup H of G . Identify an infinitesimal character χ_{inf} with (the Weyl group orbit of) an element $\lambda \in \mathfrak{h}^*$, by the Harish-Chandra homomorphism. Define $\lambda \rightarrow \bar{\lambda}$ with respect to the real form $X^*(H) \otimes \mathbb{R}$ of \mathfrak{h}^* , and write $\overline{\chi_{\text{inf}}}$ for the corresponding action on infinitesimal characters. This is well-defined, independent of all choices.

For simplicity we restrict to $GL(n, \mathbb{R})$ from now on.

Lemma 8.3 *Suppose π is an admissible representation of $GL(n, \mathbb{R})$, admitting an infinitesimal character $\chi_{\text{inf}}(\pi)$, and a central character $\chi(\pi)$. Then:*

1. $\chi_{\text{inf}}(\pi^h) = -\overline{\chi_{\text{inf}}(\pi)}$,
2. $\chi(\pi^h) = \chi(\pi)^h$,

3. Suppose $P(\mathbb{R}) = M(\mathbb{R})N(\mathbb{R})$ is a parabolic subgroup of $GL(n, \mathbb{R})$, and π_M is an admissible representation of $M(\mathbb{R})$. Then $\text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}(\pi_M^h) \simeq \text{Ind}_{P(\mathbb{R})}^{G(\mathbb{R})}(\pi_M)^h$.

Proof. The first assertion is easy if π is a minimal principal series representation. Since any irreducible representation embeds in a minimal principal series, (1) follows. Statement (2) is elementary, and (3) is an easy variant of (7.3). We leave the details to the reader. \square

Now suppose ϕ is a finite dimensional representation of $W_{\mathbb{R}}$ on a complex vector space V . Define a representation ϕ^h on the space V^h of conjugate linear functions $V \rightarrow \mathbb{C}$ by: $\phi(w)(f)(v) = f(\phi(w^{-1})v)$ ($w \in W_{\mathbb{R}}, f \in V^h, v \in V$). Choosing dual bases of V, V^h , identify $GL(V), GL(V^h)$ with $GL(n, \mathbb{C})$, to write $\phi^h = {}^t\bar{\phi}^{-1}$.

It is elementary that ϕ has a nondegenerate invariant Hermitian form if and only if $\phi \simeq \phi^h$.

Take ${}^L G = GL(n, \mathbb{C})$, so irreducible admissible representation of $GL(n, \mathbb{R})$ are parametrized by n -dimensional semisimple representations of $W_{\mathbb{R}}$. Write $\phi \rightarrow \pi(\phi)$ for this correspondence.

Lemma 8.4 *Suppose ϕ is an n -dimensional semisimple representation of $W_{\mathbb{R}}$. Then*

1. $\chi_{\text{inf}}(\phi^h) = -\overline{\chi_{\text{inf}}(\phi)}$,
2. $\chi_{\text{rad}}(\phi^h) = \chi_{\text{rad}}(\phi)^h$.

Proof. Let ${}^{\vee}H$ be the diagonal torus in $GL(n, \mathbb{C})$. As in (6.4), write $\phi(z) = z^{\lambda}\bar{z}^{\lambda'}$, so $\chi_{\text{inf}}(\phi) = \lambda$. On the other hand $\phi^h(z) = \overline{z^{\lambda}\bar{z}^{\lambda'}^{-1}}$, and it is easy to see this equals $z^{-\bar{\lambda}'}\bar{z}^{-\bar{\lambda}}$, where $\bar{\lambda}$ is complex conjugation with respect to $X_*({}^{\vee}H) \otimes \mathbb{R}$. Therefore $\chi_{\text{inf}}(\phi^h) = -\bar{\lambda}'$. Then (1) follows from the fact that, by (6.4)(d), λ is $GL(n, \mathbb{C})$ -conjugate to λ' .

The second claim comes down to the case of tori, which we leave to the reader. \square

Proof of Theorem 1.5. The equivalence of (1) and (2) follow from the preceding discussion. For (3), it is well known (and a straightforward exercise) that $\pi(\phi)$ is tempered if and only if $\phi(W_{\mathbb{R}})$ is bounded [6, 10.3(4)], which is equivalent to ϕ being unitary.

The proof of (1) is parallel to that of Theorem 1.3, using the previous two Lemmas, and our characterization of the Langlands classification in terms of infinitesimal character, radical character, and compatibility with parabolic induction. We leave the few remaining details to the reader. \square

References

- [1] J. Adams, M. van Leeuwen, A. Paul, P. Trapa, D. A. Jr. Vogan, and W.-L. Yee. Unitary dual of real reductive groups. preprint.
- [2] Jeffrey Adams. Computing global characters. In *Representation Theory and Mathematical Physics*, volume 557 of *Contemp. Math.*, pages 79–112. Amer. Math. Soc., Providence, RI, 2011.
- [3] Jeffrey Adams, Dan Barbasch, and David A. Vogan, Jr. *The Langlands classification and irreducible characters for real reductive groups*, volume 104 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1992.
- [4] Jeffrey Adams and Fokko du Cloux. Algorithms for representation theory of real reductive groups. *J. Inst. Math. Jussieu*, 8(2):209–259, 2009.
- [5] Jeffrey Adams and David A. Vogan, Jr. L -groups, projective representations, and the Langlands classification. *Amer. J. Math.*, 114(1):45–138, 1992.
- [6] A. Borel. Automorphic L -functions. In *Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2*, Proc. Sympos. Pure Math., XXXIII, pages 27–61. Amer. Math. Soc., Providence, R.I., 1979.
- [7] Michael Harris and Richard Taylor. *The geometry and cohomology of some simple Shimura varieties*, volume 151 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2001. With an appendix by Vladimir G. Berkovich.
- [8] Guy Henniart. Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p -adique. *Invent. Math.*, 139(2):439–455, 2000.

- [9] Anthony W. Knapp and David A. Vogan, Jr. *Cohomological induction and unitary representations*, volume 45 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1995.
- [10] R. P. Langlands. On the classification of irreducible representations of real algebraic groups. In *Representation theory and harmonic analysis on semisimple Lie groups*, volume 31 of *Math. Surveys Monogr.*, pages 101–170. Amer. Math. Soc., Providence, RI, 1989.
- [11] D. Shelstad. Characters and inner forms of a quasi-split group over \mathbf{R} . *Compositio Math.*, 39(1):11–45, 1979.
- [12] T. A. Springer. *Linear algebraic groups*, volume 9 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, second edition, 1998.
- [13] J. Tits. Normalisateurs de tores. I. Groupes de Coxeter étendus. *J. Algebra*, 4:96–116, 1966.
- [14] David A. Vogan, Jr. Irreducible characters of semisimple Lie groups. IV. Character-multiplicity duality. *Duke Math. J.*, 49(4):943–1073, 1982.